



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII
International Rangeland Congress

Dynamical Studies on Species Interaction and Biomass of Mixed Pasture in South-Western China

L. Q. Wan

Chinese Academy of Agricultural Sciences, China

X. L. Li

Chinese Academy of Agricultural Sciences, China

H. W. Gao

Chinese Academy of Agricultural Sciences, China

F. He

Chinese Academy of Agricultural Sciences, China

J. R. Bai

Chinese Academy of Agricultural Sciences, China

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/10-2/6>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

L. Q. Wan, X. L. Li, H. W. Gao, F. He, J. R. Bai, and F. Liu

Dynamical studies on species interaction and biomass of mixed pasture in south-western China

L. Q. Wan, X. L. Li, H. W. Gao, F. He, J. R. Bai, F. Liu

Institute of Animal Sciences of Chinese Academy of Agricultural Science, Beijing, China, E-mail: wlq@iascaas.net.cn

Key words: *Hemarthria compressa*, *Medicago sativa*, mixture, biomass, competition

Introduction The forage yields and nutritional values of mixtures with legumes and grass have been shown to be much higher than those of monocultures (Xu *et al.*, 2003). *Hemarthria compressa* and *Medicago sativa* are two dominant species in local pastures. The competitive interaction and biomass dynamics of *Hemarthria compressa* and *Medicago sativa* have not yet been studied (Moreira, 1989).

Materials and methods *Hemarthria compressa* and *Medicago sativa* (cv. Eureka, Beijing Baraern Seed Company) were grown alone and in mixtures in Hongya county of Sichuan province, China. Five treatments (three replicates per treatment, 10m² for each plot) of this study were as follows: A (100% *H. compressa*), B (75% *H. compressa* and 25% *M. sativa*), C (50% *H. compressa* and 50% *M. sativa*), D (25% *H. compressa* and 75% *M. sativa*) and E (100% *M. sativa*). As the height of vegetation comes to 35 cm, the growth of above biomass within 1m² quadrat is trimmed to a stubble of 5 cm and weighed.




Figure 1 Relative yield total (RYT) and competition ratio (CR) for mixtures of *H. compressa* and *M. sativa*. Thick lines represent RYT and thin lines (broken lines mean CR of *medicago sativa* and real lines mean CR of *H. compressa*) represent CR. The measurements were done during 2004-2005.

Results The values of relative yield total (relative to the monocultures) (RYT) ($RYT = Y_{hm}/Y_{hh} + Y_{mh}/Y_{mm}$) were higher than 1. *Hemarthria compressa* and *Medicago sativa* with same or different eco-niches performed a specific intergrowth overyielding each other in the mixtures. Otherwise, there was a significant counter-growth yield suppression between both species on the treatments. The competition ratio (CR) $CR_h = (Y_{hm}/Y_{hh} \cdot Z_{hm}) / (Y_{mh}/Y_{mm} \cdot Z_{mh})$ of *H. compressa* was higher than that of *M. sativa* on B treatment, and the reverse case on D treatment.

Conclusions There were the counter-growth and intergrowth between *H. compressa* and *M. sativa* on B treatment, and the counter-growth on C and D treatments. The competitiveness of *H. compressa* was more intensive than that of *M. sativa* on B treatment, and the reverse case on D treatment. In general, the treatment with the greatest growth was 75% *H. compressa* and 25% *M. sativa*.

Reference

Z. Xu, Y. W. Yu, S. H. Chang (2003). Effect of grazing intensity on the annual productivity and composition of mixed *Lolium perenne* and *Trifolium repens* pasture. *Acta Prataculturae Sinica*. 12(5): 31-37.